

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A61K 9/08, 39/02, 39/12, 47/10, 47/14	A1	(11) International Publication Number: WO 98/34594 (43) International Publication Date: 13 August 1998 (13.08.98)
(21) International Application Number: PCT/US98/02283 (22) International Filing Date: 3 February 1998 (03.02.98) (30) Priority Data: 60/036,900 6 February 1997 (06.02.97) US (71) Applicant (for all designated States except US): MERCK & CO., INC. [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): NG, Assunta, S. [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065 (US). HENNESSEY, John, P. [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065 (US). MANCINELLI, Ralph, J. [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065 (US). (74) Common Representative: MERCK & CO., INC.; 126 East Lincoln Avenue, Rahway, NJ 07065 (US).		(81) Designated States: CA, JP, US, European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
(54) Title: THIMEROSAL-FREE PRESERVATIVES FOR VACCINES (57) Abstract Novel combination of preservatives (methyl and propyl parabens, benzyl alcohol, and 2-phenoxyethanol) were found to pass antimicrobial testing according to USP, BP, and EP. The new preservatives were put into vaccines using L-histidine as a buffer to keep pH at 7.0. HPLC methods were developed to analyze these preservatives and their degradation products.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakhstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

TITLE OF THE INVENTION

THIMEROSAL-FREE PRESERVATIVES FOR VACCINES

CROSS-REFERENCE TO RELATED APPLICATIONS

5 Not applicable.

STATEMENT REGARDING FEDERALLY-SPONSORED R&D

Not applicable.

10 REFERENCE TO MICROFICHE APPENDIX

Not applicable.

FIELD OF THE INVENTION

15 The disclosure relates to thimerosal-free preservatives for
vaccines.

BACKGROUND OF THE INVENTION

For multidose vaccine formulations, preservatives are
required to prevent contamination of and to stabilize the composition of
20 subsequent doses after the first dose is used. The preservative must enable
the vaccine formulation to pass efficacy tests or antimicrobial challenge
tests according to the United States Pharmacopeia (USP) in the U.S.,
British Pharmacopeia (BP), and European Pharmacopeia (EP) in Europe.

Thimerosal is a commonly-used preservative in vaccines.
25 Thimerosal is a mercurial compound that is potentially toxic, and causes
allergic reaction in about sixteen percent of the population. Thimerosal is
also toxic to the environment.

It would be advantageous to find new and safer preservatives
for vaccines to replace thimerosal. In this application, we report on new
30 combinations of preservatives for vaccines: methyl and propyl parabens,
benzyl alcohol, and 2-phenoxy-ethanol. These combination preservatives
are non-toxic, yet effective. ✓

One dose of vaccine (0.5 mL) has about 1 mg paraben.
Toxicity of the parabens is relatively low, due to the ease and rapidity with
35 which the body rids itself of these drugs. The LD₅₀ of methyl paraben in
mice intraperitoneally is 1g/kg.

One dose of vaccine has about 7.5 mg benzyl alcohol. This amount is below harmful levels. Benzyl alcohol is metabolized to benzoic acid, which is conjugated with glycine in the liver, and excreted as hippuric acid. The probable lethal dose of benzyl alcohol is 0.5 - 5.0 g/kg.

5 One dose of vaccine has 2 mcL of 2-phenoxyethanol. Toxicity of 2-phenoxyethanol is low. It has been in commercial use for several decades. The presence of 2-phenoxyethanol is known in volatile naturally occurring substances, such as green tea. The acute oral LD₅₀ in rats is 1.26-2.33 mL/kg. The acute dermal LD₅₀ in rabbits is 2.0 mL/kg.

10 Due to stringent antimicrobial requirements of the various pharmacopeias, finding the right preservative for vaccine formulations is a challenge. The pH of the vaccine should be maintained at approximately pH 7. pH also has an effect on the antimicrobial effectiveness of the preservatives. Solubility of some preservatives, such as the parabens, at
15 pH 7 and at 4°C is a limiting factor. Thus, the use of combination preservatives such as methyl and propyl parabens helps to solubilize more parabens. Each paraben has its own solubility for pH 7 and 4 degrees centigrade. Using both methyl and propyl parabens together rather than separately, helps to put more paraben in solution. Methyl paraben and
20 propyl paraben work synergistically, since they exhibit differential antimicrobial activities.

The search for an effective buffer which maintains pH at pH 7 and which is safe for injectibles, is another challenge. Phosphate is the most commonly used buffer of choice for pH 7. However, phosphate buffer
25 is incompatible with many forms of aluminum hydroxide adjuvant used in vaccine formulations. Other buffers effective at this pH range may not be safe for injectibles. In this application, we report the use of L-histidine, because it is an effective buffer at pH 7, and at 20 - 35 mM final concentration is safe to use in vaccines.

30 We have developed sample preparation and high performance liquid chromatography methods for analyzing these preservatives and their degradation products in vaccines. Methods for simultaneously analyzing some of these preservatives and their degradation products are not yet present in the literature.

35

SUMMARY OF THE INVENTION

New combinations of preservatives that pass antimicrobial testing requirements for United States Pharmacopeia (USP), British Pharmacopeia (BP), and European Pharmacopeia (EP). They are: (1) 1.5% benzyl alcohol; (2) 0.225% methyl paraben sodium, 0.025% propyl paraben sodium; and 0.9% benzyl alcohol, and (3) 0.225% methyl paraben sodium, 0.025% propyl paraben sodium, and 0.375% 2-phenoxyethanol. L-histidine is used as a buffer to keep pH of vaccines neutral. A new technique for analysis of combination preservatives and their degradation products in vaccines is also provided.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1: Reversed-Phased HPLC Chromatogram of Preservative Related Components: (1) parahydroxybenzoic acid, (2) benzyl alcohol, (3) phenol, (4) benzoic acid, (5) methyl paraben, (6) benzaldehyde, and (7) propyl paraben.

Figure 2: HPLC Assay of Preservatives (Methyl Paraben, Propyl Paraben, 2-Phenoxyethanol, Benzyl Alcohol, and m-Cresol) in Vaccines.

DETAILED DESCRIPTION OF THE INVENTION

Preservatives must pass antimicrobial efficacy tests. We performed the antimicrobial tests according to United States Pharmacopeia (USP), British Pharmacopeia (BP), and European Pharmacopeia (EP). Five test organisms were used: Asperigillus niger, Candida albicans, Pseudomonas aeruginosa, Staphylococcus aureus, and Escherichia coli. We have found new combinations of preservatives that passed antimicrobial testing. Three combinations passed all antimicrobial requirements for USP, BP, and EP. They are: (1) 1.5% benzyl alcohol, (2) 0.225% methyl paraben sodium, 0.025% propyl paraben sodium, and 0.9% benzyl alcohol, and (3) 0.225% methyl paraben sodium, 0.025% propyl paraben sodium, and 0.375% 2-phenoxyethanol. Five other preservative combinations passed USP, but failed BP, and EP. They are: (4) 0.18% methyl paraben sodium, plus 0.02% propyl paraben sodium, (5) 0.9% benzyl alcohol, (6) 0.18% methyl paraben sodium, plus 0.02% propyl paraben sodium, 25 ppm formaldehyde, (7) 0.18% methyl paraben sodium,

plus 0.02% propyl paraben sodium, 0.5% benzyl alcohol, and (8) 0.27% methyl paraben sodium, plus 0.03% propyl paraben sodium.

EXAMPLE 1

5 Preparation of Vaccine Formulations

Vaccine formulations were prepared as follows. Preservatives were first prepared as concentrated stock solutions. Methyl paraben sodium is first dissolved in water at room temperature to 20% (w/v). (For example, weigh out 0.1 gm of methyl paraben sodium, and add 0.5 mL of
10 water to make a stock of 20% solution.) Propyl paraben sodium is first dissolved in water at room temperature to 2% (w/v). (For example, weight out 0.03 gm of propyl paraben sodium, and add 1.5 mL of water to make a 2% solution.) 2-Phenoxy-ethanol is first dissolved in absolute ethanol to 50% (v/v). (For example, mix 1 mL of 2-Phenoxyethanol with 1 mL of water
15 to give a 50% solution). Benzyl alcohol is used as is (v/v).

Two vaccines were studied. One is a hepatitis B vaccine, a yeast-derived recombinant hepatitis B surface antigen. The second is a combination vaccine, composed of hemophilized influenza type B, a yeast-derived recombinant hepatitis B surface antigen, diphtheria, tetanus, and
20 acellular pertussis components. Hepatitis B vaccine is Thimerosal-free, Recombivax®HB, BAP (Hepatitis B surface antigen) = 10 mcg/mL and 450 mcg aluminum hydroxide / mL.

L-histidine is used as a buffer to maintain pH 7. Buffer is added before addition of preservatives. For vaccines with parabens, L-
25 histidine stock solution (0.5 M; pH 6) is added to vaccine.

The combination vaccine was designated as AR5. AR5 is composed of Recombivax®, 10 mcg/mL HBsAg (hepatitis B surface antigen), PRP-T (ActHib), 20 mcg PRP/mL, Agglutinogens, 10 mcg/mL, 69K (Pertactin), 6 mcg/mL, Filamentous Hemagglutinen, 40 mcg/mL, LPF
30 (PT or Pertussis Toxoid), 40 mcg/mL, Diphtheria, 30 Lf/mL, and Tetanus, 10 Lf/mL, to a final concentration of 20 - 35 mM histidine. The sodium salts of parabens are at pH 9; thus using stock L-histidine buffer at pH 6 will maintain final pH at pH 7. (For example, to make 5 mL of vaccine with 0.225% methyl paraben sodium, 0.025% propyl paraben sodium, and 0.9%
35 benzyl alcohol: to a glass vial, add 4496 mcL of vaccine, 90 mcL of water,

25 mcL of 0.5 M histidine solution, pH 6, 56 mcL of 20% methyl paraben, 63 mcL of 2% propyl paraben, and 45 mcL of benzyl alcohol. Mix to dissolve.)

5

EXAMPLE 2

To make 100 mL of AR5, add 50 mL of Recombivax[®]HB, 20 mcg/mL, mixed with 2.67 mL of CLL (aluminum hydroxide; 11 mL of PRP-T (ActHib), 181.5 mcgPRP/mL; 3.48 mL of Agglutinogens, 287 mcg/mL; 2.07 mL of 69K (Pertactin), 290 mcg/mL; 5.71 mL of Filamentous Humagglutinin, 700 mcg/mL; 11.05 mL of LPF (Pertussis Toxoid), 362 mcg/mL; 0.875 mL of Diphtheria, 3430 Lf/mL mixed with 8.00 mL aluminum hydroxide; and 0.379 mL of Tetanus, 2640 Lf/mL mixed with 2.67 mL aluminum hydroxide. For vaccines preserved with benzyl alcohol alone, L-histidine solution at 0.5 M, initial pH 7.0, is added to a final concentration of 20 - 35 mM. (For example, to make 5 mL of vaccine with 1.5% benzyl alcohol: to a glass vial, add 4496 mcL of vaccine, 33 mcL of 0.5 M histidine solution, pH 7, 97 mcL of water, and 75 mcL of benzyl alcohol. Mix to dissolve.) Add preservatives slowly, a little at a time, with slow stirring, so as not to chemically or physically alter vaccine components.

15
20 Add parabens before benzyl alcohol or 2-phenoxethanol. The concentration of histidine is 20 - 35 mM. Final pH is 7. The final concentration of preservatives is as indicated in the formulation.

Preservatives are recovered by a first centrifugation to remove aluminum hydroxide adjuvant and proteins, and a second centrifugation through a 1000 molecular weight cutoff Millipore filter tube to remove all other formulation components. For example, pipette 200 mcL of vaccine with preservatives (0.225% methyl paraben sodium, 0.025% propyl paraben sodium, and 0.9% benzyl alcohol) into a 1.5 mL microcentrifuge tube. Centrifuge at maximum speed on table top microcentrifuge for 3 minutes at room temperature. Pipette out the supernatant into a clean microcentrifuge tube. Discard pellet. Pipette 40 mcL of supernatant and 160 mcL of water into a microfuge tube with 1000 molecular weight cutoff filter. Mix and centrifuge at maximum speed on table top microcentrifuge for 14 minutes at room temperature. 10 mcL of filtrate is injected into HPLC for analysis.

25
30
35

EXAMPLE 3

Preservatives such as methyl and propyl parabens, benzyl alcohol, benzoic acid, and phenol are routinely used for antimicrobial preservation in biological products. Quantitative analysis of methyl and propyl paraben by high performance liquid chromatography was popular. Quantitative analysis of methyl and propyl parabens and their degradation product, p-hydroxybenzoic acid has been carried out by thin layer chromatography (TLC) and high performance TLC. While benzyl alcohol and its degradation product, benzaldehyde, were analyzed using HPLC, other HPLC analyses of benzyl alcohol in pharmaceuticals were published. Analysis of phenol had been done by HPLC and GC.

In the course of our work, we developed a method for putting combination preservatives in biological products, facilitated by use of a buffering system for maintaining pH at 7. We also developed an efficient method of retrieving preservatives of interest for fast and accurate analysis by removing sample matrix interference. We also developed a simple HPLC method for the simultaneous separation of methyl and propyl parabens, parahydroxybenzoic acid, phenol, benzyl alcohol, benzaldehyde, and benzoic acid. Parahydroxybenzoic acid and phenol are degradation products of methyl and propyl parabens, while benzaldehyde and benzoic acid are degradation products of benzyl alcohol.

EXAMPLE 4

A Hewlett Packard HP 1090 Series HPLC consisting of autosampler, pump, and diode array detector was used. A variable wavelength detector is, however, sufficient for this work. The column was Waters- μ -Bondapak C-18, RP column (30 X 3.9 mm I.D., 10 micrometer particles). The guard column used was also Waters- μ -Bondapak. A Fisher Micro-Centrifuge Model 235A was used for centrifuging samples. A Millipore UF3 LGC WB 10,000 NMWL Filter unit was used for separating preservatives from possible sample matrix interference.

Acetonitrile was Omnisolve HPLC grade from EM Science. Benzaldehyde, benzoic acid, and phenol were from J. T. Baker. Benzyl alcohol was NF grade from A. A. Spectrum Chemical. Glacial acetic acid was Fisher Reagent ACS. L-histidine monochloride, monohydrate, was

from Spectrum Chemical Mfg. Corp. Parabenoic acid was from Sigma. Methyl paraben sodium was Nipagin M. Sodium, NF grade. Propyl paraben sodium was Nipasol M. Sodium, NF grade. Both were from Nippa Laboratories. Water was Milli-Q purified from in-house source.

5

EXAMPLE 5

Sample and Standard Preparation

The following method was used to add preservatives to the biological samples. Histidine solution at 100 mM stock, initial pH 6.0, was added as a buffer to a final concentration of 20 mM to keep the biological samples at pH 7.0 prior to addition of preservatives. Methyl and propyl parabens sodium were first dissolved in water at room temperature to 20% and 2% (w/v), respectively. Benzyl alcohol was used as is (v/v) without prior dilution. They were added to biological samples to the desired final concentrations. Standards were made fresh daily in the same manner using water instead of biological samples.

To separate the preservatives from the sample matrix for analysis, 200 ml of sample was centrifuged for 3 minutes at room temperature using the Fisher Micro-Centrifuge to remove insoluble materials. The supernatant obtained from the centrifugation was diluted with water to the desired target level, and then placed in a Millipore filter tube and centrifuged at room temperature for 14 minutes to remove additional sample matrix components. For analysis, 10 ml of filtrate was injected directly into the HPLC.

25

EXAMPLE 6

Quantitation of Preservative by HPLC

The mobile phase consisted of acetonitrile-water containing 2% (v/v) acetic acid with the following linear gradient of acetonitrile concentration: 0 min, 24%; 5 min, 24%; 9 min, 50%; 13 min, 24%. 10 ml of sample was injected. Flow rate was 2 mL/minute. Detector was set at 254 nm. Run time was 20 minutes and the assay was conducted at room temperature.

30

EXAMPLE 7

The sodium salts of methyl and propyl parabens were chosen instead of the esters, because the sodium salts are very soluble in water at room temperature. Histidine was used for these studies because histidine
5 has effective buffering capacity near pH 7 ($pK_a = 7$ at 4°C). Solutions of methyl and propyl parabens sodium have pH of about 9. To bring the pH to 7 with using only a final concentration of 20 mM histidine, an initial stock of 100 mM histidine, pH 6 was used.

After the preservatives were combined with the samples, the
10 challenge was to quantitate the concentration of the preservatives with minimal interference from the sample matrix. This was achieved by centrifugation of the sample to remove insoluble components, followed by centrifugation through a 10K MW cutoff membrane. All preservatives studied passed through the filter membrane, with a recovery of better than
15 99%.

A chromatogram of the seven preservative-related components is shown in Fig. 1. Parahydroxybenzoic acid elutes as a peak with a retention time of 2.79 minutes, benzyl alcohol at 4.18 minutes, phenol at 5.14 minutes, benzoic acid at 6.07 minutes, methyl paraben at
20 6.91 minutes, benzaldehyde at 7.95 minutes, and propyl paraben at 11.21 minutes.

Table 1 shows the linearity, intercept, and slope for standard curves of all seven compounds. The calibration graphs were constructed from two injections each of five or more concentrations. The least square
25 regression fit showed good linearity ($R\text{-square} > 0.999$) in the defined range of the standard curve for all compounds.

TABLE 1
Linearity of Compounds

	<u>Compound</u>	<u>Linear Range (mg)</u>	<u>Intercept</u>	<u>Slope</u>	<u>R-Square</u>
5	benzaldehyde	0.025 - 10.0	13.51	1844.1	0.999
	benzoic acid	0.100 - 10.0	2.29	169.9	0.999
	benzyl alcohol	0.500 - 7.5	-0.31	49.3	0.999
	methyl paraben	0.010 - 10.0	7.20	2469.0	0.999
10	parahydroxy- benzoic acid	0.100 - 2.5	36.59	2527.3	0.999
	phenol	1.000 - 5.0	-8.87	156.6	0.999
	propyl paraben	0.500 - 5.0	41.22	2195.8	0.999

Table 2 shows the reproducibility of retention times for the seven compounds. Mean values were from six replicate injections. The relative standard deviations were better than 0.3% for the seven compounds.

TABLE 2
Reproducibility of Retention Times*

	Compound	Retention Time** (minutes)	R.S.D. (%)
5	parahydroxybenzoic acid	2.79	0.25
	benzyl alcohol	4.18	0.15
10	phenol	5.14	0.26
	benzoic acid	6.07	0.28
	methyl paraben	6.91	0.29
	benzaldehyde	7.95	0.19
	propyl paraben	11.21	0.05

15

* Chromatograph in Figure 1

** Mean value of six replicates

Table 3 shows reproducibility of areas of six replicate injections. For six repeated injections in the same run, the relative standard deviations were better than 2% for all compounds. For between-day precision, seven spiked samples were analyzed in duplicate on seven separate days. The RSD (%) for methyl and propyl parabens was 5.4 and 15.1 respectively, and 8.4 for benzyl alcohol.

25

TABLE 3
Reproducibility of Areas*

	Compound	Area Units**	R.S.D. (%)
5	parahydroxybenzoic acid	3080.5	0.1
	benzyl alcohol	148.3	0.4
	phenol	167.9	0.4
10	benzoic acid	197.6	0.4
	methyl paraben	3453.0	0.1
	benzaldehyde	1103.6	1.0
	propyl paraben	359.8	2.0

15 * Chromatograph in Figure 1

** Mean value of six replicate injections

Table 4 shows the limit of detection for each compound as measured by signal-to-noise ratio of 3: 1.

20

TABLE 4
Limits of Detection and Quantitation

	<u>Compound</u>	<u>LOD (ng)</u>	<u>LOQ (ng)</u>
25	benzaldehyde	25	25
	benzoic acid	100	100
	benzyl alcohol	250	500
	methyl paraben	10	10
30	parahydroxybenzoic acid	10	100
	phenol	100	1000
	propyl paraben	25	500

35 Table 5 shows recovery studies of biological samples spiked with three different levels of preservatives. Recoveries were from 90 to

111%. As to specificity, we observed no interference from sample matrix components.

TABLE 5

5 Recovery Study of Preservatives in Biological Samples

	Compound %	Amount Added %	Recovery* %	R.S.D.*
10	methyl paraben	0.180	102	0.41
		0.225	102	0.01
		0.270	102	0.10
		0.315	104	0.16
	propyl paraben	0.020	90	0.46
15		0.025	97	0.42
		0.030	96	0.14
	benzyl alcohol	0.500	111	0.57
		1.200	111	0.49
		1.500	110	0.37
20		2.000	110	0.37

* Mean value of two repeated injections

EXAMPLE 8

25 Stability studies of the new preservative combinations in AR5 combination vaccine. AR5 is composed of Recombivax®HB, 10 mcg/mL, PRP-T (ActHib), 20 mcg/mL, Agglutinogens, 10 mcg/mL, 69K (Pertactin), 6 mcg/mL, Filamentous Hemagglutinin, 40 mcg/mL, LPF (PT or Pertussis Toxoid), 40 mcg/mL, Diphtheria, 30 Lf/mL, and Tetanus, 10 Lf/mL, and
 30 phosphate buffered saline, were done at 37°C for 7, 12, 16, and 21 days and at 4°C for 27, 57, and 96 days. There is no significant decrease in any of the preservatives concentration.

35 Table 6 shows stability studies of vaccine and phosphate buffered saline with 0.18% methyl paraben sodium, and 0.02% propyl paraben sodium. Samples for 37°C were tested for 1, 7, 12, 16 and 21 days.

Samples of 4°C were tested for 1, 27, 28, 57 and 96 days. Amount of preservatives were compared to day 1.

TABLE 6: PERCENT PRESERVATIVES COMPARED TO DAY 1
 5 0.18% METHYL PARABEN SODIUM, PLUS 0.2% PROPYL PARABEN SODIUM

		Day		1	7	12	16	21
		Batch		1	1	2	1	2
		Temperature C			37	37	37	37
Sample	Preservatives							
PBS	Methyl Paraben			100	107	96	110	106
	Propyl Paraben			100	104	88	96	99
AR5	Methyl Paraben			100	116	101	122	126
	Propyl Paraben			100	100	101	103	123
		Day		1	27	28	57	96
		Batch		1	1	2	1	2
		Temperature C			4	4	4	4
PBS	Methyl Paraben			100	103	107	107	118
	Propyl Paraben			100	96	102	100	109
AR5	Methyl Paraben			100	101	108	100	109
	Propyl Paraben			100	87	109	87	107
PBS = Phosphate Buffered Saline								
AR5 = Acthib, Recombivax, Diphtheria, Tetanus, and Pertussis								
acellular five components								

Table 7 is for 0.9% benzyl alcohol.

TABLE 7: PERCENT PRESERVATIVES COMPARED TO DAY 1
0.9% BENZYL ALCOHOL

5

		Day		1	7	12	16	21
		Batch		1	1	2	1	2
		Temperature C			37	37	37	37
<i>Samples</i>	<i>Preservatives</i>							
PBS	Benzyl Alcohol			100	120	100	141	128
AR5	Benzyl Alcohol			100	83	81	82	70
		Day		1	27	28	57	96
		Batch		1	1	2	1	2
		Temperature C			4	4	4	4
PBS	Benzyl Alcohol			100	108	121	105	101
AR5	Benzyl Alcohol			100	71	81	66	90
PBS = Phosphate Buffered Saline								
AR5 = Acthib, Recombivax, Diphtheria, Tetanus, and Pertussis								
acellular 5 components								

Table 8 is for 0.6% phenoxyethanol.

TABLE 8: PERCENT PRESERVATIVES COMPARED TO DAY 1
0.6% PHENOXYETHANOL

5

		Day		1	7	12	16	21
		Batch		1	1	2	1	2
		Temperature C			37	37	37	37
<i>Sample</i>	<i>Preservatives</i>							
PBS	Phenoxyethanol			100	100	97	107	118
AR5	Phenoxyethanol			100	88	81	106	87
		Day		1	27	28	57	96
		Batch		1	1	2	1	2
		Temperature C			4	4	4	4
PBS	Phenoxyethanol			100	96	115	101	130
AR5	Phenoxyethanol			100	73	87	79	115
PBS = Phosphate Buffered Saline								
AR5 = Acthib, Recombivax, Diphtheria, Tetanus, and Pertussis								
acellular 5 components								

Table 9 for 0.18% methyl paraben sodium, 0.02% propyl paraben sodium, and 0.25% phenoxyethanol.

TABLE 9: PERCENT PRESERVATIVES COMPARED TO DAY 1
0.18% METHYL PARABEN SODIUM, 0.2% PROPYL PARABEN
SODIUM, AND 0.25% 2-PHENOXYETHANOL

		Day		1	7	12	16	21
		Batch		1	1	2	1	2
		Temperature C			37	37	37	37
Sample s	Preservatives							
PBS	Methyl Paraben			100	103	98	118	122
	Propyl Paraben			100	96	92	104	115
	2-Phenoxyethanol			100	111	95	122	115
AR5	Methyl Paraben			100	102	101	118	136
	Propyl Paraben			100	94	99	108	127
	2-Phenoxyethanol			100	100	97	107	119
		Day		1	27	28	57	96
		Batch		1	1	2	1	2
		Temperature C			4	4	4	4
PBS	Methyl Paraben			100	102	108	102	121
	Propyl Paraben			100	92	105	91	112
	2-Phenoxyethanol			100	108	106	105	100
AR5	Methyl Paraben			100	100	104	99	95
	Propyl Paraben			100	93	108	91	95
	2-Phenoxyethanol			100	99	102	95	77
PBS = Phosphate Buffered Saline								
AR5 = Acthib, Recombivax, Diphtheria, Tetanus, and Pertussis								
acellular 5 components								

Table 10 for 0.18% methyl paraben sodium, 0.2% propyl paraben sodium, and 25 ppm formaldehyde.

5 TABLE 10: PERCENT PRESERVATIVES COMPARED TO DAY 1
0.18% METHYL PARABEN SODIUM, 0.2% PROPYL PARABEN
SODIUM, AND 25 PPM FORMALDEHYDE

		Day		1	7	12	16	21
		Batch		1	1	2	1	2
		Temperature C			37	37	37	37
<i>Sample</i>	<i>Preservatives</i>							
PBS	Methyl Paraben			100	104	140	118	
	Propyl Paraben			100	91	128	94	
AR5	Methyl Paraben			100	105	102	118	119
	Propyl Paraben			100	96	101	109	114
		Day		1	27	28	57	96
		Batch		1	1	2	1	2
		Temperature C			4	4	4	4
PBS	Methyl Paraben			100	103	105	103	105
	Propyl Paraben			100	86	98	87	89
AR5	Methyl Paraben			100	103	106	102	107
	Propyl Paraben			100	97	107	95	107
PBS = Phosphate Buffered Saline								
AR5 = Acthib, Recombivax, Diphtheria, Tetanus, and Pertussis acellular 5 components								

5 TABLE 11: PERCENT PRESERVATIVES COMPARED TO DAY 1
 0.18% METHYL PARABEN SODIUM, 0.2% PROPYL PARABEN
 SODIUM, AND 0.5% BENZYL ALCOHOL

		Day		1	7	12	16	21
		Batch		1	1	2	1	2
		Temperature C			37	37	37	37
Sample s	Preservatives							
PBS	Methyl Paraben			100	104	104	108	112
	Propyl Paraben			100	91	94	89	101
	Benzyl Alcohol			100	107	103	105	103
AR5	Methyl Paraben			100	105	101	126	123
	Propyl Paraben			100	96	99	115	118
	Benzyl Alcohol			100	96	98	102	109
		Day		1	27	28	57	96
		Batch		1	1	2	1	2
		Temperature C			4	4	4	4
PBS	Methyl Paraben			100	105	105	106	115
	Propyl Paraben			100	89	99	89	93
	Benzyl Alcohol			100	105	100	106	99
AR5	Methyl Paraben			100	98	105	100	104
	Propyl Paraben			100	89	105	93	91
	Benzyl Alcohol			100	92	101	93	89
PBS = Phosphate Buffered Saline								
AR5 = Acthib, Recombivax, Diphtheria, Tetanus, and Pertussis								
acellular 5 components								

WHAT IS CLAIMED:

1. A method of stabilizing vaccines comprising mixing stock vaccine solutions with nonmercurial preservatives and L-histidine
5 buffer.
2. The method of Claim 1 wherein the preservatives are selected from the group consisting of approximately 1.5% benzyl alcohol; a mixture of approximately 0.225% methyl paraben sodium, approximately
10 0.025% propyl paraben sodium, and approximately 0.9% benzyl alcohol; and a mixture of approximately 0.225% methyl paraben sodium, approximately 0.025% propyl paraben sodium, and approximately 0.375% 2-phenoxyethanol.
- 15 3. Vaccines prepared by the method of Claim 1.
4. Vaccines prepared by the method of Claim 2.
5. A method of determining stability of the vaccines of
20 Claim 3 comprising HPLC analysis of preservatives and degradation products of preservatives.
6. A method of determining stability of the vaccines of Claim 4 comprising HPLC analysis of preservatives and degradation
25 products of preservatives.




FIG.1

2/2




FIG.2

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US98/02283

A. CLASSIFICATION OF SUBJECT MATTER IPC(6) :A61K 9/08, 39/02, 39/12, 47/10, 47/14 US CL :424/201.1, 202.1, 203.1, 227.1, 247.1, 245.1, 256.1, 514/21, 785 According to International Patent Classification (IPC) or to both national classification and IPC		
B. FIELDS SEARCHED Minimum documentation searched (classification system followed by classification symbols) U.S. : 424/201.1, 202.1, 203.1, 227.1, 247.1, 245.1, 256.1, 514/21, 785 Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched The Merck Index, 11th Edition Electronic data base consulted during the international search (name of data base and, where practicable, search terms used) APS, JPO, EPO, STN: MEDLINE, EMBASE, BIOSIS, PATOSWO vaccine, benzyl alcohol, paraben, phenoxyethanol, HPLC, preservative, degradation product, not mercury, histidine		
C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	LOWE, I, et al., The Antimicrobial Activity of Phenoxyethanol in Vaccines. Let. Appl. Microbiol. 1994. Vol. 18. pages 115-116, see the abstract.	1-4
Y	KNECZKE, M. Determination of Pilocarpine, Physostigmine, its Degredation Product Reserine and Preservatives by High Performance Liquid Chromatography. 1980. Vol. 198. pages 529-533, see entire document.	5, 6
Y	MONATH, T.P. Stability of Yellow Fever Vaccine. Develop. Biol. Standard. 1996. Vol. 87, pages 219-225, especially page 221.	1-4
<input checked="" type="checkbox"/> Further documents are listed in the continuation of Box C. <input type="checkbox"/> See patent family annex.		
* "A" "B" "L" "O" "P"	Special categories of cited documents: document defining the general state of the art which is not considered to be of particular relevance earlier document published on or after the international filing date document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) document referring to an oral disclosure, use, exhibition or other means document published prior to the international filing date but later than the priority date claimed	*T* *X* *Y* *A* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art document member of the same patent family
Date of the actual completion of the international search 08 APRIL 1998		Date of mailing of the international search report 02 JUN 1998
Name and mailing address of the ISA/US Commissioner of Patents and Trademarks Box PCT Washington, D.C. 20231 Facsimile No. (703) 305-3230		Authorized officer MARY TUNG Telephone No. (703) 308-0196

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US98/02283

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	CAMERON, J. Preservative Systems Compatible with DPT-Polio (Salk) and TABTD-Polio (Salk) Vaccines. Develop. Biol. Standard. 1974. Vol. 24. pages 155-165, especially page 156.	1-4
Y	EP 0,750,907 A2 (AMERICAN CYANAMID COMPANY) 02 January 1997, page 6, lines 19-21.	1-4
Y, P	US 5,603,933 A (DWYER et al.) 18 February 1997, column 11, lines 45-49.	1-4
Y, P	US 5,672,350 A (PARKER et al.) 30 September 1997, column 10, lines 60-67.	1-4